



doi: <https://doi.org/10.20546/ijcrar.2023.1105.002>

Antibiotic Property of Edible and Non-Edible Wild Mushrooms from Menengai Crater in Kenya

Paul Njenga Waithaka^{1*} and Githaiga Benson Muriuki²

¹School of Pure and Applied Sciences, Kirinyaga University, P.O. Box, 143, Kerugoya

²Department of Biological Sciences Egerton University, P. O. Box, 536, Njoro, Kenya

*Corresponding author

Abstract

Antibiotic resistance is a global challenge of our times. To avert the problem, search for alternative antibiotics from natural sources is inevitable. This study sought to extract antibiotics from edible and non-edible wild mushrooms from Menengai crater. The mushroom samples were collected in new khaki bags from Menengai crater and transported to the laboratory. The samples were dried under room temperature ($25\pm 2^{\circ}\text{C}$). Extraction of the crude extracts was carried out using ethyl acetate, methanol, ethanol and acetone. Antibiotic bioassay was done using Kirby Bauer disc diffusion technique against *Enterococcus faecalis* (ATCC 29212), *Staphylococcus aureus* (ATCC 25923), *Pseudomonas aeruginosa* (ATCC 27853), *Klebsiella pneumoniae* (ATCC 70063), *Aspergillus flavus* (ATCC 9170), *Aspergillus fumigatus* (96918) and *Candida albicans* (10231). The yield of crude extract obtained using ethyl acetate, methanol, ethanol and acetone varied significantly ($P=0.009$). Likewise, the total phenolic content varied significantly among the extracts ($P=0.000105$). However, the zones of inhibition did not vary significantly among the bacterial ($P=0.98$) and fungal pathogens ($P=0.61$). The selected mushroom samples produced bioactive compounds that inhibited growth of the selected pathogenic microorganisms. There is need to characterize the active compounds in the crude extracts obtained from the mushroom samples.

Article Info

Received: 08 March 2023

Accepted: 20 April 2023

Available Online: 20 May 2023

Keywords

Antibiotic, Extracts, Menengai, Mushroom, Solvents, fatty acid content.

Introduction

Human dietary requirements are changing with changing lifestyle all over the world (New and Zin, 2020). The demand for alternative healthy sources of nutrients and medicine is on the increase. This can be partly attributed to increase in antibiotic resistance (Shen *et al.*, 2017). Microbial products are a major constituent of new drug molecules. Researchers are searching for novel antimicrobial agents of biological origin including fungi (Waithaka *et al.*, 2017).

More than 2000 species of mushroom exist in nature of which 25 are used as food while a few are commercially cultivated. They are of importance nutritionally because of their organoleptic merit, medicinal properties and economic significance (Uddin *et al.*, 2015).

Mushrooms are of high nutritional value (Rezaeian and Pourianfar, 2016). They are rich in protein and fibre while their fat content is low but with incomparable fatty acid content. Besides, they provide vitamin B complex, C, D and E (Mohamed and Farghaly, 2014). Mushrooms

have extensively been used in traditional medicine in curing intestinal disorders, bleeding, high blood pressure and various bacterial infection (Mirfat *et al.*, 2014). The Metabolites from mushroom have been shown to have antimicrobial, antioxidant and anti-inflammatory properties (Alves *et al.*, 2012). Besides, antibiotics from mushrooms have anti-obesity, anti-diabetic and anti-cancer properties (Shikongo *et al.*, 2013). Their antibiotic properties are attributed to procession of bioactive compounds such as terpenoids, flavonoids, tannins, alkaloids, tannins, alkaloids and polysaccharides (Christiane *et al.*, 2016).

Ethyl acetate, ethanol, methanol and water have previously been used in extracting antibiotics from mushrooms (Thillaimahanan *et al.*, 2016). Water is usually used because of its high polarity. The extraction capacity of water can be increased though increasing its temperature (Aina *et al.*, 2012). However, high temperature may lead to degradation of antimicrobials that are heat sensitive (Celik *et al.*, 2014). This study sought to investigate the antibiotic properties of wild mushrooms from Menengai crater.

Materials and Methods

Sample collection

Two edible (*Lactarius kabansus* and *Boletus edulis*) and two non-edible (*Coprinus comatus* and *Ganoderma lucidum*) wild mushrooms were collected from Menengai crater and transported to the Department of Biological Sciences Laboratory in Egerton University. The samples were identified using morphological characteristics and identification keys (Khan and Tania, 2012).

Extraction of crude extracts

The samples were sliced into thin strips and dried for 7d under room temperature ($25\pm 2^\circ\text{C}$). The samples were separately ground to powder using an electrical grinder (Siebtechnik steel pulverizer 2, 376, GmbH). Briefly, 100g of the powder was separately mixed with 100mL of ethyl acetate, methanol, ethanol and acetone and shaken using an orbital shaker at 150 rpm for 72h (Sagar *et al.*, 2022).

The extracts were centrifuged at 3000 rpm for 15 min. Filtration was carried out using Whatman No.1 filter paper. The extracts were concentrated using a rotary evaporator at 50°C , freeze dried and weights determined (Hussein *et al.*, 2018).

Test pathogenic Microorganisms

The bacterial test pathogens were *Enterococcus faecalis* (ATCC 29212), *Staphylococcus aureus* (ATCC 25923), *Pseudomonas aeruginosa* (ATCC 27853) and *Klebsiella pneumoniae* (ATCC 70063). The fungi used as test organisms were: *Aspergillus flavus* (ATCC 9170), *Aspergillus fumigates* (96918) and *Candida albicans* (10231).

Determination of total phenolic content

Total phenolic content in each mushroom extract was determined using the Folin and Ciocalteu (FC) reagent method (Padmavathy *et al.*, 2014). Briefly, 40 μl of each sample was diluted to 200 μl using distilled water and mixed with 200 μl of Folin and Ciocalteu's phenol reagent, diluted in the ratio of 1:9 mL using distilled water. After 6 min, 200 μl of 7.5% sodium carbonate was added to the mixture and adjusted to 2mL with distilled water and stored in the dark for 60 min. The absorbance was measured at 725 nm using a spectrophotometer (SpectronicR) (Deepalakshmi and Mirunalini, 2014).

Antibiotic bioassay

Antimicrobial activity of the crude extracts was carried out using Kirby Bauer disc diffusion bioassay. Briefly, each extract was dissolved in dimethyl sulfoxide (DMSO) to a concentration of 200 mg/mL and sterilized by passage through a ChromafilXtra PTFE-45 filter. Sterile Mueller Hinton Agar, was dispensed into 90mm Petri dishes. Whatmann No:1 paper was used to prepare 6mm discs which were sterilized using a hot air oven at 150°C for 1h (Woldegiorgis *et al.*, 2014). Test microorganism cultures incubated for 24h were adjusted to a cell density of 1.5×10^8 CFU/mL as per 0.5 McFarland Standard.

A micropipette was used to dispense 100 μl on Mueller Hinton agar and spread using an L shaped glass rod. The Discs were aseptically dipped into 10 μl of each extract. The discs were dried under room temperature ($25\pm 2^\circ\text{C}$) before placement at certain points within the Petri dish.

Distilled water was used as negative control while ampicillin 10 μg for bacteria and nystatin 10 μg for fungi disc were used as positive control. The Petri dishes were stored at 4°C for 2 h to allow absorption of the extracts and then incubated at 37°C for 24 h for bacterial and 5d at 25°C for fungal pathogens. The diameter of zones of inhibition were measured in millimeters.

Statistical analyses

All the statistical analyses were carried out using SPSS version 25.0 software. All values are expressed as mean + SD of three parallel measurements.

Results and Discussion

Yield of extracts

The percentage yield in ethyl acetate varied from 4.7 ± 0.01 to 6.5 ± 0.02 , methanol (2.5 ± 0.03 - 3.4 ± 0.02), ethanol (3.0 ± 0.03 - 3.5 ± 0.03) and acetone (4.1 ± 0.02 - 4.7 ± 0.03) (Table 1). The crude extracts yield obtained from the mushroom samples using the selected extracts varied significantly ($P=0.009$).

Total phenolic content

The total phenolic content of the crude extracts obtained using ethyl acetate ranged from 75 ± 0.01 to 87 ± 0.03 mg GAE/ 100g, methanol (47 ± 0.01 - 69 ± 0.02 mg GAE/ 100g), ethanol (45 ± 0.02 - 58 ± 0.01 mg GAE/ 100g) and acetone (77 ± 0.02 - 86 ± 0.02 mg GAE/ 100g) (Table 2). The total phenolic content in the extracts varied significantly ($P= 0.000105$).

Growth inhibition of bacterial pathogens

Growth inhibition of *Enterococcus faecalis* varied from 13 ± 0.01 - 19 ± 0.02 mm, *Staphylococcus aureus* (13 ± 0.01 - 19 ± 0.01 mm), *Pseudomonas aeruginosa* (10 ± 0.02 - 15 ± 0.03 mm) and *Klebsiella pneumoniae* (10 ± 0.01 - 15 ± 0.03 mm) (Table 3). However, the zones of inhibition did not vary significantly ($P=0.98$) among the test bacterial pathogens.

Growth inhibition of fungal pathogens

The zones of inhibition in *Aspergillus flavus* ranged from 10 ± 0.01 to 13 ± 0.03 mm, *Aspergillus fumigatus* (10 ± 0.03 - 15 ± 0.01 mm) and *Candida albicans* (14 ± 0.03 - 18 ± 0.01 mm). The zones of inhibition obtained did not vary significantly ($P=0.61$) between *Aspergillus flavus*, *Aspergillus fumigatus* and *Candida albicans*. However, the zones of inhibition obtained in *Candida albicans* were higher than in *Aspergillus flavus* and *Aspergillus fumigatus*.

The crude extracts obtained using ethyl acetate was higher than in methanol, ethanol and acetone. This may be attributed to differences in polarity of the solvents (Chowdhury *et al.*, 2015). A previous study on Antioxidant profile of four selected wild edible mushrooms in Nigeria obtained similar results (Ejelonu *et al.*, 2013). However, the % yield of the crude extracts in the present study was higher than in a study carried out by Özcan and Ertan, (2018). The difference could be attributed to differences in the environmental conditions in which the samples were growing (Pathirage and Yunman, 2011).

In addition, non-edible (*Coprinus comatus* and *Ganoderma lucidum*) mushrooms produced higher crude extracts than the edible (*Lactarius kabansus* and *Boletus edulis*) mushrooms. This concurred with a study on antimicrobial activity of crude ethanolic extracts from some medicinal mushrooms carried out by Lukáš *et al.*, (2016). Differences in biosynthesis pathways between the edible and non-edible mushrooms may have contributed to the results (Gan *et al.*, 2013).

The results on total phenolic content obtained in the current study were higher than in a previous study carried out by Sun *et al.*, (2014). The difference could be attributed to the geographical conditions in which the sample mushrooms were obtained (Gan *et al.*, 2013).

According to Sagar *et al.*, (2015), the temperature of the area in which mushrooms grow significantly determines their level of crude extracts. In addition, Asri *et al.*, (2019) maintained that the type of substrate on which mushrooms grow determines their phenolic content.

The zones of inhibition of the selected bacterial pathogens by crude extracts from the sampled mushroom obtained in the current study were larger than those obtained in a previous study by Chowdhury *et al.*, (2015). Differences in the bioactive compounds present in the crude extracts could be a contributing factor (Geethangili *et al.*, 2013). The zones of inhibition obtained in the Gram positive bacteria were bigger than those from Gram negative bacteria. Roy *et al.*, (2016) maintained that differences in the structure of the cellular components in bacterial pathogens result in larger zones of inhibition in Gram positive than in Gram negative bacteria.

Table.1 Yield (%) of the crude extracts from the mushroom samples

Mushroom	Solvent			
	Ethyl acetate	Methanol	Ethanol	Acetone
<i>Lactarius kabansus</i>	5.7±0.02	2.7±0.01	3.0±0.03	4.3±0.01
<i>Boletus edulis</i>	4.7±0.01	2.5±0.03	3.2±0.01	4.1±0.02
<i>Coprinus comatus</i>	6.0±0.03	3.4±0.02	3.5±0.03	4.7±0.03
<i>Ganoderma lucidum</i>	6.5±0.02	3.2±0.01	3.4±0.02	4.6±0.01

Table.2 Total phenolic content (mg GAE/ 100 g) of mushrooms extracted using different solvents.

Mushroom extract	Total phenolic content			
	Ethyl acetate	Methanol	Ethanol	Acetone
<i>Lactarius kabansus</i>	78±0.01	59±0.02	47±0.03	77±0.02
<i>Boletus edulis</i>	75±0.01	47±0.01	45±0.02	79±0.01
<i>Coprinus comatus</i>	90±0.02	67±0.01	57±0.01	86±0.02
<i>Ganoderma lucidum</i>	87±0.03	69±0.02	58±0.01	85±0.01

Table.3 Zones of inhibition (mm) of the test bacterial pathogens by the crude extracts from the selected mushroom samples

Pathogen	Zone of inhibition			
	Ethyl acetate	Methanol	Ethanol	Acetone
<i>Enterococcus faecalis</i>				
<i>Lactarius kabansus</i>	14±0.01	14±0.02	13±0.03	15±0.02
<i>Boletus edulis</i>	15±0.01	13±0.01	15±0.02	13±0.01
<i>Coprinus comatus</i>	18±0.02	17±0.01	17±0.01	17±0.02
<i>Ganoderma lucidum</i>	17±0.03	19±0.02	18±0.01	19±0.01
<i>Staphylococcus aureus</i>				
<i>Lactarius kabansus</i>	13±0.01	15±0.02	14±0.03	13±0.03
<i>Boletus edulis</i>	14±0.02	13±0.02	15±0.02	14±0.02
<i>Coprinus comatus</i>	17±0.02	18±0.01	15±0.01	16±0.02
<i>Ganoderma lucidum</i>	16±0.03	19±0.01	17±0.02	18±0.02
<i>Pseudomonas aeruginosa</i>				
<i>Lactarius kabansus</i>	10±0.02	12±0.01	11±0.03	10±0.02
<i>Boletus edulis</i>	11±0.01	10±0.02	10±0.02	12±0.01
<i>Coprinus comatus</i>	13±0.02	14±0.01	15±0.03	14±0.01
<i>Ganoderma lucidum</i>	15±0.01	14±0.01	13±0.01	13±0.02
<i>Klebsiella pneumoniae</i>				
<i>Lactarius kabansus</i>	12±0.03	10±0.02	11±0.01	10±0.03
<i>Boletus edulis</i>	10±0.01	11±0.02	11±0.01	12±0.01
<i>Coprinus comatus</i>	14±0.02	15±0.01	15±0.03	14±0.02
<i>Ganoderma lucidum</i>	13±0.01	14±0.01	13±0.02	15±0.01

Table.4 Zones of inhibition (mm) of the test fungal pathogens by the crude extracts from the selected mushroom samples

Pathogen	Zone of inhibition			
	Ethyl acetate	Methanol	Ethanol	Acetone
<i>Aspergillus flavus</i>				
<i>Lactarius kabansus</i>	11±0.02	13±0.03	12±0.01	13±0.03
<i>Boletus edulis</i>	12±0.01	10±0.01	13±0.03	11±0.01
<i>Coprinus comatus</i>	10±0.01	11±0.02	11±0.03	10±0.02
<i>Ganoderma lucidum</i>	13±0.03	12±0.01	12±0.02	12±0.01
<i>Aspergillus fumigatus</i>				
<i>Lactarius kabansus</i>	13±0.02	10±0.03	12±0.02	11±0.01
<i>Boletus edulis</i>	14±0.03	12±0.02	11±0.01	13±0.02
<i>Coprinus comatus</i>	15±0.01	11±0.02	13±0.02	12±0.01
<i>Ganoderma lucidum</i>	14±0.01	13±0.01	11±0.01	14±0.01
<i>Candida albicans</i>				
<i>Lactarius kabansus</i>	18±0.01	15±0.02	17±0.03	18±0.01
<i>Boletus edulis</i>	16±0.02	14±0.03	18±0.01	17±0.02
<i>Coprinus comatus</i>	16±0.01	16±0.01	17±0.02	16±0.02
<i>Ganoderma lucidum</i>	17±0.03	16±0.02	16±0.02	18±0.01

Candida albicans presented larger zones of inhibition than *Aspergillus flavus* and *Aspergillus fumigatus*. Nedelkoska *et al.*, (2013) asserted that moulds are more resistant to crude extracts than unicellular fungi. Besides, the ability of moulds to produce resistant spores accords them the ability to resist inhibition by crude extracts (Tsongai *et al.*, 2016).

The results of the present study concurred with previous studies carried out elsewhere (Hleba *et al.*, 2016; Reid *et al.*, 2016; Roy *et al.*, 2016). This may be attributed to production of similar active ingredients by the sample mushrooms (Krishnaveni and Manikandan, 2014).

Ethyl acetate, methanol, ethanol and acetone extracted bioactive materials from the selected mushroom samples. Ethyl acetate was the best solvent for recovering crude extracts from mushroom samples. Polarity of the solvent is key in the extraction process. The crude extracts from the mushroom samples contained phenolic compounds. In addition, the crude extracts inhibited growth of the selected bacterial and fungal pathogens

Recommendations

There is need to characterize the crude extracts from the selected mushroom samples. The crude extracts obtained in this study need to be tested against other microbial pathogens. Extraction of crude extracts from other mushroom samples need to be carried out.

References

- Aina, D. A., Jonathan, S. G., Olawuyi, O. J., Ojelabi, D. O. and Durowoju, B. M. (2012). Antioxidant, antimicrobial and phytochemical properties of alcoholic extracts of *Cantharellus cibarius*– a Nigerian mushroom. *New York Science Journal*; 5(10):114-120.
- Alves, M. J., Ferreira, I. C., Dias, J., Teixeira, V. and Martins, A. (2012). A review on antimicrobial activity of mushrooms (Basidiomycetes) extracts and isolated compounds. *Plant med*; 78: 1707-1718.
- Asri, R.M., Yahya, H., Rehan, M. M. and Yahya, H. N. (2019). Antibacterial Properties of Ethanolic Extract of Mushrooms Sold in Malaysian Local Market. *East African Scholars Journal of Agriculture and Life Sciences*; 2 (11):516-523.
- Celik, G. Y., Onbasli, D., Altinsoy, B. and Alli, H. (2014). *In vitro* antimicrobial and antioxidant properties of *Ganoderma lucidum* extracts grown in Turkey. *European Journal of Medicinal Plants*; 4(6):709-722.
- Chowdhury, M. M. H., Kubra, K., and Ahmed, S. R. (2015). Screening of antimicrobial, antioxidant properties and bioactive compounds of some edible mushrooms cultivated in Bangladesh. *Annals of Clinical Microbiology and Antimicrobials*; 14(1):8-15.

- Cristiane, U. J., Eliana, F., Margô, G. O. and Karnikowski, J. K. (2016). Antimicrobial properties of the mushroom *Agaricus blazei* – integrative review. *Brazilian Journal of Pharmacognosy*; 26:780-786.
- Deepalakshmi, K., and Mirunalini, S. (2014). Assessment of in vitro antioxidant and antimicrobial properties of cultivated *Pleurotus ostreatus*: An edible mushroom. *Free Radicals and Antioxidants*; 4(2):123-129.
- Ejelonu, O. C., Akinmoladun, A. C., Elekofehinti, O. O. and Olaleye, M. T. (2013). Antioxidant profile of four selected wild edible mushrooms in Nigeria. *Journal of Chemical Pharmaceutical Research*; 7:286–45.
- Gan, C. H., Narul, A. B. and Asmah, R. (2013). Antioxidant analysis of different types of edible mushrooms (*Agaricus bisporus* and *Agaricus brasiliensis*). *Int. Food Research Journal*; 20(3):1095-1102.
- Geethangili, M., Rao, Y. K. and Tzeng, Y. M. (2013). Development and validation of HPLC-DAD separation method for determination of bioactive anthrocon medicinal mushroom *Antrodia camphorata*. *International Journal of Applied Science Engineering*; 11(2):195-201.
- Hleba, L., Kompas, M., Hutková, J., Rajtar, M., Petrová, J., Cubon, J. and Káčániová, M. (2016). Antimicrobial activity of crude ethanolic extracts from some medicinal mushrooms. *The Journal of Microbiology, Biotechnology and Food Sciences*; 5:60-69.
- Hussein, A., Mohammed, E. and Hamid, E. (2018). Antibacterial activity of alcoholic and aqueous extracts of *A. bisporus* against food borne bacterial pathogens. *Al-Nahrain Journal of Science*; 21:111-114.
- Khan, A. and Tania, M. (2012). Nutritional and medicinal importance of *Pleurotus* mushrooms: An overview. *Food Reviews International*; 28(3):313-329.
- Krishnaveni, M. and Manikandan, M. (2014). Antimicrobial activity of mushrooms. *Research Journal of Pharmacy and Technology*; 7:399-400.
- Lukáš, H., Matúš, K., Jana, H., Martin, R. and Jana, P., (2016). Antimicrobial activity of crude ethanolic extracts from some medicinal mushrooms. *Journal of Microbiology and Biotechnology of Food Sciences*; 5: 60-63.
- Mirfat, A. H. S., Noorlidah, A., and Vikineswary, S. (2014). Antimicrobial activities of split gill mushroom *Schizophyllum commune* Fr. *American Journal of Research Communication*; 2(7): 113- 124.
- Mohamed, E. M. and Farghaly, F. A. (2014). Bioactive compounds of fresh and dried *Pleurotus ostreatus* mushroom. *International Journal of Biotechnology for Wellness Industries*; 3(1):4-14.
- Nedelkoska, D. N., Panjcevska, N.A., Amedi, H, Veleska, D., Ivanova, E. and Karadelev, M. (2013). Screening of antibacterial and antifungal activities of selected Macedonian wild mushrooms. *Matica Srpska. Journal of Natural Sciences*; 124:333-340.
- Özcan, Ö. and Ertan, F. (2018). Beta-glucan Content, Antioxidant and Antimicrobial Activities of Some Edible Mushroom Species. *Food Science and Technology* 6(2): 47-55.
- Padmavathy, M., Sumathy, R., Manikandan, N. and Kumuthakalavalli, R. (2014). Antimicrobial activity of mushrooms against skin infection causing pathogens. *Research in Biotechnology*; 5(2):22-26.
- Pathirage, K. P. and Yunman, L. (2011). Mushrooms as a functional food mediator in Preventing and ameliorating diabetes. *Functional Food*; 4:161-71.
- Reid, T., Kashangura, C., Chidewe, C., Benhura, M. A. and Mduluza, T. (2016). Antibacterial properties of wild edible and non-edible mushrooms found in Zimbabwe. *African Journal of Microbiology Research*; 10(26): 977-984.
- Rezaeian, S. and Pourianfar, H. R. (2016). Antimicrobial properties of the button mushroom, *Agaricus bisporus*: A mini-review. *International Journal of Advanced Research*; 4(1):426-429.
- Roy, D. N., Azad, A. K., Sultana, F. and Anisuzzaman, A. S. (2016). In-vitro antimicrobial activity of ethyl acetate extract of two common edible mushrooms. *Journal of Pharmacology*; 5:79-82.
- Sagar, A., Aghnihotri, S. and Thakur, K. (2015). Studies on antibacterial activity of wild edible mushroom – *Lactarius vellereus*. *Indian Journal of Mushrooms*; 33(1):12-18.
- Sagar, A., Kumar, S., Kamal, S. and Rana, J. (2022). Studies on evaluation of antibacterial activities of some cultivated mushrooms against human pathogenic bacteria. *Mushroom Research*; 31(1): 81-91.
- Shen, H. S., Shao, S., Chen, J. C., and Zhou, T. (2017). Antimicrobials from mushrooms for assuring food safety. *Comprehensive Reviews in Food Science and Food Safety*; 16(2):316-329.

- Shikongo, L. T., Chimwamurombe, P. M., Lotfy, H. R. and Kandawa-Schulz, M. (2013). Antimicrobial screening of crude extracts from the indigenous *Ganoderma lucidum* mushrooms in Namibia. *African Journal of Microbiology Research*; 7(40):4812-4816.
- Sun, L., Bai, X. and Zhuang, Y. (2014). Effect of different cooking methods on total phenolic contents and antioxidant activities of four *Boletus* mushrooms. *J. Food Sci. Tech.* 51:3362-3368.
- Thillaimaharan, K. A., Sharmila, K., Thangaraju, T., Karthick, M. and Kalaiselvam, M. (2016). Studies on antimicrobial and antioxidant properties of oyster mushroom *Pleurotus florida*. *International Journal of Pharmaceutical Sciences Research*; 4:1540-1545.
- Tsungai, R., Chenjerayi, K., Catherine, C., Mudadi, A. B. and Takafira, M. (2016). Antibacterial properties of wild edible and non-edible mushrooms found in Zimbabwe. *African Journal of Microbiology Research*; 4: 977-984.
- Uddin, G. M., Hossain, S., Monirul Islam, M., Asaduzzaman, M., Bulbul, J., and Amin, R. (2015). Evaluation of antimicrobial, antioxidant and cytotoxic property of *Pleurotus ostreatus* mushroom. *International Research Journal of Biological Sciences*; 4(1):29-33.
- Waithaka, P. N., Gathuru, E. M., Githaiga, B. M., and Onkoba, K. M. (2017). Antimicrobial activity of mushroom (*Agaricus bisporus*) and fungal (*Trametes gibbosa*) extracts from mushrooms and fungi of Egerton main Campus, Njoro Kenya. *Journal Biomedical Sciences*; 6 (3): 1-6.
- Woldegiorgis, A. Z., Abate, D., Haki, G. D. and Ziegler, G. R. (2014). Antioxidant property of edible mushrooms collected from Ethiopia. *Food Chemistry*; 157:30-36.

How to cite this article:

Paul Njenga Waithaka and Githaiga Benson Muriuki. 2023. Antibiotic Property of Edible and Non-Edible Wild Mushrooms from Menengai Crater in Kenya. *Int.J.Curr.Res.Aca.Rev.* 11(05), 8-14.
doi: <https://doi.org/10.20546/ijcrar.2023.1105.002>