



doi: <https://doi.org/10.20546/ijcrar.2018.601.006>

## Simple Renal Cyst: An Independent Risk Factor of Hypertension

**S. S. Gupta, Manish Bansal\* and Varun Gupta**

*Department of Medicine, FH Medical College, Tundla, India*

*\*Corresponding author*

### Abstract

The detection of asymptomatic renal cysts is increasing with advancement of imaging techniques and availability. The aim of our study was to evaluate the presence of simple cysts with prehypertension and hypertension in our population. All consecutive adult patients aged > 25 years who attended the medicine OPD of FHMC and Hospital were enrolled in study. Detailed medical history and physician examination was done in all the patients. Abdominal ultrasonography and biochemical parameters were done. All patients who had history or evidence of structural or functional kidney disease were excluded. A total of 5000 patients were included in study and divided into three groups: normotension (2810), prehypertension (1488) and hypertension (702) groups. There were significant difference in age, sex, family history of hypertension, regular exercise, smoking habits, BMI, systolic and diastolic blood pressure, fasting plasma glucose, total Cholesterol, HDL Cholesterol, Creatinine, estimated GFR in all three groups. Both number (> 2) and size (>2) of SRCs were associated with prehypertension and hypertension independently. The presence of SRCs is independent risk factor and SRCs size >2 cm and number > are significant determinations if prehypertension and hypertension.

### Article Info

Accepted: 31 December 2017  
Available Online: 20 January 2018

### Keywords

Asymptomatic renal cysts,  
Imaging techniques,  
Hypertension

### Introduction

Simple renal cysts are discrete lesions within kidney that are typically cortical, extending outside the parenchyma. They are oval or circular in shape and have a distinct outline with enclosed liquid or semisolid fluid structures in kidney, originating secondary to genetic or acquired processes. They are quite common space occupying lesions of the kidney. Polycystic kidney disease is well known entity associated with hypertension, hematuria, flank pain and recurrent urinary tract infections with

ultimate renal function impairment.<sup>9</sup> Simple renal cysts (SRCs) are more common and considered acquired and benign. They are generally unilateral, solitary and asymptomatic except when complications occur due to infection, hemorrhage or rupture leading to formation of complex cyst.<sup>2,3,14</sup> Imaging modalities such as ultrasound, CT Scan and Magnetic resonance Imaging (MRI) are quite frequently used in the evaluation of abdominal ailments. This has led to detection of asymptomatic renal cysts that may pass unnoticed from eyes of clinicians. Initially simple renal cysts were overlooked but now

clinical importance has grown. There are no clinical data or reports are available on relation with simple renal cysts and hypertension or prehypertension in our population. The objective of our study was to evaluate the patients of SRCs and its association with hypertension and prehypertension.

### Materials and Methods

In a cross section study 5000 adults patients aged more than 25 years were enrolled in F.H. Medical College and Hospital, Firozabad (India), Medicine OPD between July 2015 to Dec 2016. Patients with history or family history of polycystic kidney disease, Medullary sponge kidney, renal disease (congenital or acquired) diabetes mellitus and pregnancy were excluded. A written and informed consent were taken from all patients which were included in this study and approved by institutional ethics committee.

Detailed medical history, family history of chronic disease and personal history were taken. Smoking status was defined as one pack of cigarette per day and patients were divided into current, past and non-smokers. Regular exercise was defined as >30 minutes for at least thrice a week. Anthropometric parameters were measured. Blood pressure was recorded by mercury sphygmomanometer. Two readings of both systolic and diastolic pressure at 15 minutes interval were taken at right arm supine position. Average of two readings were classified as normotensive (<120/80 mmHg); prehypertensive (blood pressure 120-139/80-89 mmHg) and hypertensive (Blood pressure 130/90 mmHg) according to seventh Joint National Committee (JNC VII) criteria<sup>16</sup>. Abdominal ultrasound examination was done after 8 hour overnight fast by experienced ultrasonologist. The diagnostic criteria for simple renal cysts were echolucent, round or oval shape, thin wall, well defined sharply demarcated possession wall without calcification and Doppler signals from within the cyst as well as sound wave amplification behind the cystic renal mass which was not suggestive of malignancy.<sup>11</sup> Fasting blood glucose, serum creatinine and lipid profile were measured in each patient. Glomerular filtration rate (EGFR) was estimated using the abbreviated equation from the modification of diet in Renal Disease Study.<sup>10</sup>

### Statistical Analysis

Data analysis was carried out using an SPSS software version 15 (SPSS, Chicago, IL). The X<sup>2</sup> test was used for comparison of categorical variables among groups.

Multiple linear regression was used to test effect of SRCs on systolic and diastolic pressure respectively.

### Results and Discussion

A total of 5000 patients were enrolled and divided into three groups: normotensive (n=2810) Prehypertensive (n=1488) and hypertension (n=702) groups. It was observed that there was significant difference in clinical parameters like age, sex, body mass index, family history of hypertension, regular exercise, smoking habit, systolic and diastolic blood pressure (Table 1) among three groups. The biochemical parameters like fasting plasma glucose, total cholesterol, triglyceride, HDL Cholesterol, creatinine and estimated glomerular filtration rate also shows significant difference among three groups (Table 2). Similarly number and size of renal cysts also significant difference among all three groups. (Table 3). Adjusted odds ratio (OR) and 95% confidence interval of clinical variables for the risk of prehypertension and hypertension were shown in table 4 and they are already considered as independently associated factors of prehypertension and hypertension. Renal cyst size >2 cm and number >2 were also significantly associated with both prehypertension and hypertensive groups. (Table 5)

Our study shows that presence of SRCs is independent high risk factor for prehypertension and hypertension irrespective of other metabolic, renal and cardiovascular risk factors. The size of SRCs > 2cm and number >2 also increase the magnitude of risk. Other results also show that the presence of SRCs is associated with raised systolic and diastolic blood pressure both. Pederson JF<sup>12</sup> *et al.*, observed significant association between simple renal cysts and arterial blood pressure. Similar observations were also seen in certain cross sectional observational studies.<sup>6,7,8</sup> In our study clinical parameters like age, male gender, obesity, family history of hypertension, smoking and physical activity were significantly associated with prehypertension and hypertension, although they are well known risk actors of atherosclerosis.

The mechanism of increase in blood pressure by SRCs is not yet clear. The possible explanation is that elevation of blood pressure is related to activation of rennin-angiotension-aldosterone system which can be stimulated by SRCs. Cyst expansion many cause renal ischemia leading to increased release of rennin.<sup>4,5,12,13</sup> There are certain studies that have demonstrated decreased plasma rennin activity and normalization of blood pressure after surgical removal or decompression of cysts.

**Table.1** Clinical parameters among patients with normotension, prehypertension and hypertension

S. No.	Variable	Normotension (n = 2810)	Prehyperetension (n = 1488)	Hypertension (n= 702)	P Value
1	Age(Years)	44.6 + 10.2	50.8 + 11.6	57.2 + 11.8	0.011
2	Male Gender	1516 (54%)	893 (60%)	435 (62%)	0.002
3	Body Mass Index (>23 kg/m <sup>2</sup> )	843 (30%)	759 (51%)	386 (55%)	0.01
4	Family history of hypertension	135 (5%)	198 (14%)	168 (24%)	0.015
5	Regular exercise	218 (9.2%)	97 (6.5%)	43 (6.1%)	0.05
6	Smoking	275 (9.8%)	187 (12.6%)	99 (14.2%)	0.02
7	Systolic Blood Pressure (mmHg)	112 + 9.0	127 + 7	158 + 15	0.01
8	Diastolic Blood Pressure (mmHg)	72 + 7	78 + 8	94 + 12	0.01

**Table.2** Biochemical parameters among patients with normotension, prehypertension and hypertension

S. No.	Variable	Normotension (n = 2810)	Prehyperetension (n = 1488)	Hypertension (n= 702)	P Value
1	Fasting Plasma glucose (mg/dl)	88.4 + 9.4	90.6 + 10.2	94.2 + 10.4	0.02
2	Total Cholesterol (mg/dl)	128 + 32.4	206 + 37.4	209 + 38.6	0.04
3	Triglyceride (mg/dl)	94 + 18	112 + 31	128 + 37	0.02
4	HDL Cholesterol (mg/dl)	52 + 12.4	46 + 12.7	40.8. + 11.6	.016
5.	eGFR (ml/minute/ 1.73m <sup>2</sup> )	98 + 17.6	92.6 + 15.4	86.2 + 21.2	.001
6	Creatinine (mg/dl)	0.80 + 0.08	0.86 + 0.18	1.2 + 0.72	.02

eGFR – estimated glomerular Filtration rate,  
 HDL – High Density Lipoprotein

**Table.3** Simple Renal Cysts among patients with normotension, prehypertension and hypertension

S. No.	Variable	Normotension (n = 2810)	Prehyperetension (n = 1488)	Hypertension (n= 702)	P Value
1	Presence of SRCs	167 (6%)	296 (16%)	168 (24%)	.016
2	Number >2 <2	42 (1.5%) 126 (4.5%)	111 (6%) 185 (10%)	70 (10%) 98 (14%)	0.01
3	Size >2 cm <2 cm	56 (2%) 112 (4%)	130 (7%) 166 (9%)	63 (9%) 105 (15%)	0.01

**Table.4** The adjusted odd ratio and 95% confidence interval of clinical variables on the risk of prehypertension and hypertension by multinomial logistic regression analysis

Sl. No.	Variables	Prehypertension Vs Normotension	P Value	Hypertension Vs Normotension	P Value
1	Age > 60 yrs	4.31 (3.5– 5.71)	0.001	18.62(16.14-24.16)	0.001
2	Male Vs Female	2.22 (1.90-2.43)	0.02	2.16(1.98_2.38)	0.02
3	BMI >23 Kg/m <sup>2</sup>	2.67 (2.31-2.96)	0.03	2.76(1.98-2.38)	0.01
4	Family History of Hypertension	1.38 (1.12–1.68)	0.01	1.54(1.25-1.90)	0.022
5	Regular Exercise	1.44 (1.16–1.69)	0.01	1.72(1.52-1.94)	0.01
6	Smoking	1.35 (1.16–1.64)	0.42	1.65(1.25-1.91)	0.03
7	Total Cholesterol (>200 mg/dl)	1.66 (1.39–1.92)	0.038	1.97(1.56-2.25)	0.02
8	Low HDL Cholesterol (mg/dl)	0.91 (0.62–1.13)	0.68	1.04(0.84-1.25)	0.05
9.	eGFR	0.94 (0.90–1.10)	0.55	1.60(1.40-2.10)	0.02

**Table.5** The adjusted odd ratio and 95% confidence interval of simple renal cysts on the risk of prehypertension and hypertension by multinomial logistic regression analysis

Sl. No.	Variables	Prehypertension Vs Normotension	P Value	Hypertension Vs Normotension	P Value
1	Simple Renal Cyst (Yes or No)	1.41 (1.26– 1.73)	0.02	1.86 (1.68– 2.15)	0..12
2	Simple Renal Cyst Number >2	1.97 (1.77 – 2.27)	0.001	2.33 (2.12 – 2.78)	0.001
3	Simple Renal Cyst Size >2cm	2.06 (1.74 – 2.54)	0.02	2.74 (2.20 – 3.41)	0.01

This supports the view that SRCs can cause hypertension.<sup>10,15</sup> The possible mechanism may be that even small intraparenchymal cysts can create high internal hydrostatic pressure causing renal ischemia by compressing surrounding renal tissue.<sup>13</sup>

In our study we have observed significant association between SRCs and prehypertension and hypertension. The limitation of our study are that we have no measured plasma renin activity, serum renin level and effect of surgical removal of cysts to support the hypothesis of role of increased renin levels in patients of SRCs. Therefore follow-up in larger population survey are required in future to address these issues.

With the advancement of imaging techniques, the diagnosis of simple cysts is increased. The presence of simple renal cysts (SRCs) should not be overlooked. It is an important and independent determinant of prehypertension and hypertension. SRCs >2 in number or > 2 cm in size are frequently associated with hypertension and prehypertension. Patients with SRCs must be monitored closely for blood pressure in routine clinical practice.

## References

1. Ahallal Y, Khallouk a, Tazi MF, Tazi E, Elfassi MJ, Faith MH. Remission of hypertension after treatment of giant simple renal cyst: a case report. *Cases J* 2009; 2:9152
2. Caglioti A, Esposito c, Fuiano G el al. Prevalence of symptoms in patients with simple renal cysts. *BMJ* 1993; 306: 430-431.
3. Chang CC, Kuo JY, Chan WL, chan KK, chan LS/ Prevalence and clinical characteristics of simple renal cysts. *J Chin Med Assc* 2007; 70:486-491.
4. Chapman AB, Johnson A, Gabow PA, et al., the rennin-angiotensin-aldosterone system and autosomal dominant polycystic kidney disease. *N. Eng J Med* 1990; 323:1091-1096.
5. Grantham JJ, Acquired cystic kidney disease, *Kidney Int* 1991; 40:143-152.
6. Hong S, Lim JH, Jeong IG, choe J, Kim CS, Hong JH. What association exists between hypertension and simple renal cyst in a screened polulation? *J Hum Hypertens* 2013 Mar 7. Doi: 10, 1038/jhh.2013.12.
7. Lee CT, Yang YC, Wu JS et al., Multiple and large simple renal cysts are associated with prehypertension and hypertension. *Kidney Int.* 2013; 83:924-30
8. Lee YJ, Kim MS, Cho S, Kim SR, Association between simple renal cysts and development of hypertension inhealthy middle aged men. *J Hypertens* 2012; 30: 700-4
9. Lippert MC. Renal Cystic Disease. In L Gillenwater Jt, Howards SS, Mitchell ME (eds) *Adult an Pediatric Urology*, 4<sup>th</sup> edn, Lippincott Williams and Wilkins: Philadelphia, 2002, pp 858-59
10. Levey AS, Bosch JP, Lewis JB, et al., A more accurate method to estimate glomerular filtration rate from serum creatinine: a new prediction equation. Modification of diet in Renal Disease Study Group. *Ann Intern Med* 1999; 130: 461-470.
11. Nahm AM, Ritz E. The simple renal cyst. *Nephrol Dial Transplant* 2000; 15: 1702-4
12. Pedersen JF, Emamian SA, Neilsen MB. Significant association between simple renal cysts and arterial blood pressure. *Br J Urol* 1997; 79: 688-791.
13. Pedersen JF, Emamian SA, Neilsen MB. Simple renal cyst: relation to age and arterial blood pressure. *Br J Radiol* 1993; 66: 581-584.
14. Terada N, Arai Y, Kinyukava N, Terai A. The 10 year natural history of simple renal cysts. *Urology* 2008; 71: 7-12.
15. Zerem E, Imamovic G, Omerovic S. simple renal cysts and arterial hypertension: does their evacuation decrease the blood pressure/ *J Hypertens* 2009; 40: 143-152
16. Chabonian AV, Bakris GL, Black HR et al., The Seventh report of the Joint National Committee on prevention, detection, evaluation and treatment of high blood pressure: The JNC 7<sup>th</sup> report *JAMA* 2003;289:2560-2572

### How to cite this article:

Gupta S. S., Manish Bansal and Varun Gupta. 2018. Simple Renal Cyst: An Independent Risk Factor of Hypertension. *Int.J.Curr.Res.Aca.Rev.* 6(1), 35-39. doi: <https://doi.org/10.20546/ijcrar.2018.601.006>